

A binary infinitesimal form of Teichmüller metric

GUOWU YAO

May 14, 2006

Abstract

Let S be a Riemann surface of analytic finite type or the unit disk in the complex plane. Let $[\mu]$ denote the Teichmüller equivalence classes of Beltrami differentials μ . We apply the Fundamental Inequalities to obtain a binary infinitesimal form of Teichmüller metric. Using this form, we define “*angle*” between two geodesics originating from a point and conjecture that the sum of the angles of a triangle in $T(S)$ should be less than π if S is of analytic finite type. As a consequence, the well-known necessary condition for two geodesics coinciding is derived immediately.

REFERENCES

- [1] A. F. Beardon, *The Geometry of Discrete Groups*, Springer-Verlag, New York, 1983.
- [2] H. Busemann, *The Geometry of Geodesics*, Academic Press, New York, 1955.
- [3] C. J. Earle, *The Teichmüller distance is differentiable*, Duke Math. J. 44 (1977), 389-397.
- [4] C. J. Earle, I. Kra and S. L. Krushkal, *Holomorphic motions and Teichmüller spaces*, Trans. Amer. Math. Soc. 343 (1994), 927-948.
- [5] C. J. Earle and Z. Li, *Isometrically embedded polydisks in infinite-dimensional Teichmüller spaces*, J. Geom. Anal. 9 (1999), 51-71.
- [6] F. P. Gardiner and N. Lakic, *Quasiconformal Teichmüller Theory*, Amer. Math. Soc. Providence, RI, 2000.

2000 *Mathematics Subject Classification*. Primary 32G15, 30C75; Secondary 30C62.

Key words and phrases. Teichmüller space, geodesic disk, Teichmüller disk, fundamental inequality.

The author was supported by the National Natural Science Foundation of China (Grant No. 10401036) and a Foundation for the Author of National Excellent Doctoral Dissertation (Grant No. 200518) of PR China.

- [7] S. Kravetz, *On the geometry of Teichmüller spaces and the structure of their modular groups*, Ann. Acad. Sci. Fenn. Ser. A I Math. 278 (1959), 1-35.
- [8] Z. Li, *Non-uniqueness of geodesics in infinite-dimensional Teichmüller spaces*, Complex Var. Theory Appl. 16 (1991), 261-272.
- [9] Z. Li, *Non-uniqueness of geodesics in infinite dimensional Teichmüller spaces (II)*, Ann. Acad. Sci. Fenn. Ser. A I Math. 18 (1993), 355-367.
- [10] O. Lehto and K. I. Virtanen, *Quasiconformal mappings in the plane*, Springer-Verlag, Berlin and New York, 1973.
- [11] M. R. Linch, *On metrics in Teichmüller spaces*, Ph.D. thesis, Columbia University, 1971.
- [12] M. Rees, *Teichmüller distance for analytically finite surfaces is C^2* , Proc. London Math. Soc. 85 (2002), 686-716.
- [13] H. L. Royden, *Automorphisms and isometries of Teichmüller space*, in: *Advances in the Theory of Riemann Surfaces* (Ann. Math. Studies no. 66), 369-383, Princeton Univ. Press, Princeton, New Jersey, 1971.
- [14] H. Tanigawa, *Holomorphic families of geodesic discs in infinite dimensional Teichmüller spaces*, Nagoya Math. J. 127 (1992), 117-128.

Department of Mathematical Sciences
Tsinghua University
Beijing, 100084, People's Republic of China
e-mail: gwyao@math.tsinghua.edu.cn